

On Tiling Rectangles via the Frobenius Number

J.L. Ramírez Alfonsín,
Université Pierre et Marie Curie, Paris 6,
Combinatoire et Optimisation
Case 189 - 4 Place Jussieu Paris 75252 Cedex 05, France

February 2, 2008

Abstract

In this paper, we give some sufficient conditions for a n -dimensional rectangle to be tiled with a set of bricks. These conditions are obtained by using the so-called Frobenius number.

1 Introduction

Let $a_1, \dots, a_n$ be positive integers. We denote by $A = (a_1 \times \dots \times a_n)$ the n -dimensional rectangle of sides a_i , that is, $A = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid 0 \leq x_i \leq a_i, i = 1, \dots, n\}$. A n -dimensional rectangle A is said to be *tiled* with *bricks* (i.e., small n -dimensional rectangles) $B_1, \dots, B_k$ if A can be filled entirely with copies of B_i , $1 \leq i \leq k$ (rotations allowed). It is known [4, 7] that rectangle $A = (a_1 \times a_2)$ can be tiled with $(x_1 \times x_2)$ if and only if x_1 divides a_1 or a_2 , x_2 divides a_1 or a_2 and if $x_1 x_2$ divides one side of A then the other side can be expressed as a nonnegative integer combination of x_1 and x_2 . In 1995, Fricke [5] gave the following characterization when $n = 2$ (see also [3] for a n -dimensional generalization with $k = 2$).

Theorem 1.1 [5] *Let a_1, a_2, x, y be positive integers with $\text{g.c.d.}(x, y) = 1$. Then, $(a_1 \times a_2)$ can be tiled with $(x \times x)$ and $(y \times y)$ if and only if either a_1 and a_2 are both multiple of x or a_1 and a_2 are both multiple of y or one of the numbers a_1, a_2 is a multiple of both x and y and the other can be expressed as a nonnegative integer combination of x and y .*

Let us consider the following natural question.

Question 1.2 *Does there exist a function $C = C(x_1, x_2, y_1, y_2)$ such that if $a_1, a_2 \geq C$ then $(a_1 \times a_2)$ can be tiled with $(x_1 \times x_2)$ and $(y_1 \times y_2)$ for some positive integers x_1, x_2, y_1 and y_2 ?*

An algebraic result due to Barnes [1] seems to show the existence of such C . However, Barnes's method does not give an explicit lower bound for C .

The special case when $x_1 = 4, x_2 = 6, y_1 = 5$ and $y_2 = 7$ was posed in the 1991 William Mowell Putnam Examination (Problem B-3). In this case, Klosinski *et. al.* [8] gave a lower bound of C . Their method was based on knowledge of the *Frobenius number*. The *Frobenius number*, denoted by $g(s_1, \dots, s_n)$, of a set of relatively prime positive integers $s_1, \dots, s_n$, is defined as the largest integer that is not representable as a nonnegative integer combination of $s_1, \dots, s_n$. It is well known that

$$g(s_1, s_2) = s_1 s_2 - s_1 - s_2. \quad (1)$$

However, to find $g(s_1, \dots, s_n)$, for general n , is a difficult problem from the computational point of view; we refer the reader to [11] for a detailed discussion on the Frobenius number. Klosinski *et. al.* used equation (1), with particular integers s_1 and s_2 , to show that $(a_1 \times a_2)$ can be tiled with (4×6) and (5×7) if $a_1, a_2 \geq 2214$.

In this paper, we will use the Frobenius number in a more general way to show that a n -dimensional rectangle A can be tiled with some set of bricks if the sides of A are larger than a certain function (see Theorem 2.1). We use then Theorem 2.1 to obtain the following result.

Corollary 1.3 *Let $a_1, a_2, p, q, r, s \geq 2$ be integers with $r < s$, $\text{g.c.d.}(qs, qr, rs) = 1$ and $\text{g.c.d.}(p, r) = \text{g.c.d.}(p, s) = \text{g.c.d.}(r, s) = 1$. Then, $(a_1 \times a_2)$ can be tiled with $(p \times q)$ and $(r \times s)$ if*

$$a_1, a_2 \geq \max\{2qrs - (qs + qr + rs) + 1, (p-1)(s-1), (r-1)(s-1)\}.$$

In the case when $p = 6, q = 4, r = 5$ and $s = 7$, Corollary 1.3 implies that $(a_1 \times a_2)$ can be tiled with (4×6) and (5×7) if $a_1, a_2 \geq \max\{198, 20, 24\} = 198$, improving the lower bound given in [8]. We remark that this lower bound is not optimal. In [9], Narayan and Schwenk showed that, in this particular case, it is enough to have $a_1, a_2 \geq 33$. However, their tiling constructions allow rotations of both bricks (and tilings with more complicated patterns) which is not the case of Corollary 1.3.

We shall also use Theorem 2.1 to prove the following result concerning tilings of squares.

Theorem 1.4 *Let $1 < p_1 < \dots < p_{n+1}$ be prime integers. Then, $(\underbrace{a \times \dots \times a}_n)$ can be tiled with $(\underbrace{p_1 \times \dots \times p_1}_n), \dots, (\underbrace{p_{n+1} \times \dots \times p_{n+1}}_n)$ if*

$$a > n \prod_{i=1}^{n+1} p_i - \sum_{i=1}^{n+1} \frac{p_1 \dots p_{n+1}}{p_i}.$$

We finally improve the lower bound given in Theorem 1.4 in some special cases.

Theorem 1.5 *Let $p > 4$ be an odd integer with $3 \nmid p$ and let a be a positive integer. Then, $(a \times a)$ can be tiled with $(2 \times 2), (3 \times 3)$ and $(p \times p)$ if $a \geq 3p + 2$.*

Moreover, $(a \times a)$ can be tiled with (2×2) , (3×3) and (5×5) if and only if $a \neq 1, 7$ and with (2×2) , (3×3) and (7×7) if and only if $a \neq 1, 5, 11$.

A collection of some unpublished work, due to D.A. Klarner, in relation with Theorem 1.5 can be found in [10].

2 Tilings

We need to introduce some notation and definitions. Let $B = \{b_1, \dots, b_k\}$ where b_i are positive integers. We will write $g.c.d.(B)$ instead of $g.c.d.(b_1, \dots, b_k)$ and $g(B)$ instead of $g(b_1, \dots, b_k)$. Let x_j^i be a positive integer for each $j = 1, \dots, n$ and each $i = 1, \dots, n+1$. Let $\{i_1 < \dots < i_{k+1}\} \subseteq \{1, \dots, n+1\}$, $1 \leq k \leq n$. We define the set

$$X_k(i_1, \dots, i_{k+1}) = \left\{ \frac{x_k^{i_1} \cdots x_k^{i_{k+1}}}{x_k^{i_{k+1}}}, \dots, \frac{x_k^{i_1} \cdots x_k^{i_{k+1}}}{x_k^{i_1}} \right\}$$

Note that if $k = n$ then $X_n = \{x_n^1 \cdots x_n^n, x_n^1 \cdots x_n^{n-1} x_n^{n+1}, \dots, x_n^2 \cdots x_n^{n+1}\}$ is unique.

We denote by $(r; X)$ the rectangle obtained from $X = (x_1 \times \dots \times x_n)$ by sticking together r copies of X along the n^{th} -axis, that is, $(r; X) = (x_1 \times \dots \times x_{n-1} \times r x_n)$. Finally, we denote by $\bar{X}$ the $(n-1)$ -dimensional rectangle obtained from $X = (x_1 \times \dots \times x_n)$ by setting $x_n = 0$, that is, $\bar{X} = (x_1 \times \dots \times x_{n-1})$.

Theorem 2.1 *Let $x_j^i \geq 2$ be an integer for each $j = 1, \dots, n$ and each $i = 1, \dots, n+1$, $n \geq 1$. Suppose that $g.c.d.(X_k(i_1, \dots, i_{k+1})) = 1$ for any $\{i_1 < \dots < i_{k+1}\} \subseteq \{1, \dots, n+1\}$, $1 \leq k \leq n$ and let*

$$g_n = \max_{1 \leq k \leq n} \{g(X_k(i_1, \dots, i_{k+1})) | \{i_1, \dots, i_{k+1}\} \subseteq \{1, \dots, n+1\}\}$$

Then, $(a_1 \times \dots \times a_n)$ can be tiled with bricks $X^i = (x_1^i \times \dots \times x_n^i)$, $i = 1, \dots, n+1$ if $a_j > g_n$ for all $1 \leq j \leq n$.

Proof. We shall use induction on n . For $n = 1$ we have that $g.c.d.(x_1^1, x_1^2) = 1$ and thus $g_1 = g(x_1^1, x_1^2) = x_1^1 x_1^2 - x_1^1 - x_1^2$. By definition of the Frobenius number, any integer $a_1 > g_1$ is of the form $a_1 = u x_1^1 + v x_1^2$ where u, v are nonnegative integers. Thus, the 1-dimensional rectangle (a_1) (that is, the interval $[0, a_1]$) can be tiled by sticking together $(u; X^1)$ (that is, the interval $[0, u x_1^1]$) and $(v; X^2)$ (that is, the interval $[0, v x_1^2]$).

We suppose that it is true for $n = m - 1 \geq 1$ and let x_j^i be a positive integer for each $j = 1, \dots, m$ and each $i = 1, \dots, m+1$ with $g.c.d.(X_k(i_1, \dots, i_{k+1})) = 1$ for any $\{i_1 < \dots < i_{k+1}\} \subseteq \{1, \dots, m+1\}$, $1 \leq k \leq m$ and let $X^i = (x_1^i \times \dots \times x_m^i)$, $i = 1, \dots, m+1$ and $a_j > g_{m+1}$ for all $1 \leq j \leq m$.

By induction $(a_1 \times \dots \times a_{m-1})$ can be tiled with bricks $\bar{X}^{i_1}, \dots, \bar{X}^{i_m}$ for any $\{i_1 < \dots < i_m\} \subseteq \{1, \dots, m+1\}$ since $a_j \geq g_{m-1}$ for all $1 \leq j \leq m-1$.

We claim that $(a_1 \times \cdots \times a_{m-1} \times x_m^{i_1} \cdots x_m^{i_m})$ can be tiled with bricks $X^{i_1}, \dots, X^{i_m}$ for any $\{i_1 < \cdots < i_m\} \subset \{1, \dots, m+1\}$.

Indeed, if we consider the rectangle $(a_1 \times \cdots \times a_{m-1})$ embedded in $\mathbb{R}^m$ with $x_m = 0$ then by replacing each brick $\bar{X}^{i_j}$ used in the tiling of $(a_1 \times \cdots \times a_{m-1})$ by $(\frac{x_m^{i_1} \cdots x_m^{i_m}}{x_m^{i_j}}; X^{i_j})$ we obtain a tiling of $(a_1 \times \cdots \times a_{m-1} \times x_m^{i_1} \cdots x_m^{i_m})$ with bricks $X^{i_1}, \dots, X^{i_m}$.

Now, since $a_m > g_m$ then $a_m = w_{m+1}(\frac{x_m^1 \cdots x_m^{m+1}}{x_m^{m+1}}) + \cdots + w_1(\frac{x_m^1 \cdots x_m^{m+1}}{x_m^1})$ where each w_i is a nonnegative integer. By the above claim, $Y^j = (a_1 \times \cdots \times a_{m-1} \times \frac{x_m^1 \cdots x_m^{m+1}}{x_m^j})$ can be tiled with bricks $\{X^1, \dots, X^{m+1}\} \setminus X^j$ for each $j = 1, \dots, m+1$. Thus, $(a_1 \times \cdots \times a_{m-1} \times a_m)$ can be tiled with $X^1, \dots, X^{m+1}$ by sticking together bricks $(w_1; Y^1), \dots, (w_{m+1}; Y^{m+1})$ along the m^{th} -axis. $\square$

An old result due to Brauer and Shockley [2] states that if $s_1, \dots, s_n$ are positive integers with $\text{g.c.d.}(s_1, \dots, s_n) = 1$ and $\text{g.c.d.}(s_1, \dots, s_{n-1}) = d$ then

$$g(s_1, \dots, s_n) = dg\left(\frac{s_1}{d}, \dots, \frac{s_{n-1}}{d}, s_n\right) + (d-1)s_n \quad (2)$$

Remark 2.2 If $s_1, \dots, s_n$ are positive integers with $\text{g.c.d.}(s_1, \dots, s_n) = 1$ and if s_n is representable as a nonnegative integer combination of the other s_i then $g(s_1, \dots, s_n) = g(s_1, \dots, s_{n-1})$.

We shall use equation (2) and Remark 2.2 to prove Corollary 1.3.

Proof of Corollary 1.3. We consider Theorem 2.1 with $n = 2$, $(x_1^1 \times x_2^1) = (p \times q)$, $(x_1^2 \times x_2^2) = (r \times s)$ and $(x_1^3 \times x_2^3) = (s \times r)$. Then, $(a_1 \times a_2)$ can be tiled with $(p \times q)$ and $(r \times s)$ if

$$a_2, a_2 > g_2 = \max\{g(p, r), g(p, s), g(r, s), g(qs, qr, sr)\}.$$

By equation (2), $g(qs, qr, rs) = qg(s, r, rs) + (q-1)rs$ and, by Remark 2.2, $g(qs, qr, rs) = qg(s, r) + (q-1)rs = q(sr - s - r) + qrs - rs = 2qrs - (qr + qs + rs)$. The result follows by noticing that $g(p, r) < g(p, s)$ since $r < s$. $\square$

In order to prove Theorem 2.1, we need the following lemma.

Lemma 2.3 Let $1 < p_1 < \cdots < p_{n+1}$ be prime integers, $n \geq 1$. Then,

$$(a) \quad g\left(\frac{p_1 \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_1 \cdots p_{n+1}}{p_1}\right) = n \prod_{i=1}^{n+1} p_i - \sum_{i=1}^{n+1} \frac{p_1 \cdots p_{n+1}}{p_i}.$$

(b) For each $k = 1, \dots, n$,

$$g\left(\frac{p_{n-k+1} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-k+1} \cdots p_{n+1}}{p_{n-k+1}}\right) \geq g\left(\frac{p_{i_1} \cdots p_{i_{k+1}}}{p_{i_{k+1}}}, \dots, \frac{p_{i_1} \cdots p_{i_{k+1}}}{p_{i_1}}\right)$$

for all $\{i_1 < \cdots < i_{k+1}\} \subseteq \{1, \dots, n+1\}$ with equality if and only if $\{i_1, \dots, i_{n+1}\} = \{1, \dots, n+1\}$.

Proof. (a) By induction on n . For $n = 1$ we have that $g(p_1, p_2) = p_1 p_2 - p_1 - p_2$ finding equation (1). We suppose that it is true for $n = m$. We assume that $n = m + 1$, by equation (2), we have

$$g\left(\frac{p_1 \cdots p_{m+1}}{p_{m+1}}, \dots, \frac{p_1 \cdots p_{m+1}}{p_1}\right) = p_1 g\left(\frac{p_2 \cdots p_{m+1}}{p_{m+1}}, \dots, \frac{p_2 \cdots p_{m+1}}{p_2}, \frac{p_1 \cdots p_{m+1}}{p_1}\right) + (p_1 - 1) \frac{p_1 \cdots p_{m+1}}{p_1}$$

Since $\frac{p_1 \cdots p_{m+1}}{p_1} = p_2 \frac{p_2 \cdots p_{m+1}}{p_2}$ then, by Remark 2.2,

$$g\left(\frac{p_2 \cdots p_{m+1}}{p_{m+1}}, \dots, \frac{p_2 \cdots p_{m+1}}{p_2}, \frac{p_1 \cdots p_{m+1}}{p_1}\right) = g\left(\frac{p_2 \cdots p_{m+1}}{p_{m+1}}, \dots, \frac{p_2 \cdots p_{m+1}}{p_1}\right)$$

and thus

$$\begin{aligned} g\left(\frac{p_1 \cdots p_{m+1}}{p_{m+1}}, \dots, \frac{p_1 \cdots p_{m+1}}{p_1}\right) &= p_1 g\left(\frac{p_2 \cdots p_{m+1}}{p_{m+1}}, \dots, \frac{p_2 \cdots p_{m+1}}{p_1}\right) + (p_1 \cdots p_{m+1}) - \frac{p_1 \cdots p_{m+1}}{p_1} \\ &= (m-1) \prod_{i=1}^{m+1} p_i - \sum_{i=2}^{m+1} \frac{p_1 \cdots p_{m+1}}{p_i} + \prod_{i=1}^{m+1} p_i - \frac{p_1 \cdots p_{m+1}}{p_1} \\ &= p_1 \left((m-1) \prod_{i=2}^{m+1} p_i - \sum_{i=2}^{m+1} \frac{p_2 \cdots p_{m+1}}{p_i} \right) + \prod_{i=1}^{m+1} p_i - \frac{p_1 \cdots p_{m+1}}{p_1} \\ &= m \prod_{i=1}^{m+1} p_i - \sum_{i=1}^{m+1} \frac{p_1 \cdots p_{m+1}}{p_i} \end{aligned}$$

(b) By induction on k . For $k = 1$ we have $g(p_{i_1}, p_{i_2}) = (p_{i_1} - 1)(p_{i_2} - 1) \leq (p_n - 1)(p_{n+1} - 1) = g(p_n, p_{n+1})$ since $p_{i_1} \leq p_n$ and $p_{i_2} \leq p_{n+1}$ for all $\{i_1 < i_2\} \subset \{1, \dots, n+1\}$. We suppose that it is true for $k = m$ and assume that $k = m + 1 \leq n$. By equation (2) we have

$$\begin{aligned} g\left(\frac{p_{n-m} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-m} \cdots p_{n+1}}{p_{n-m}}\right) &= p_{n-m} g\left(\frac{p_{n-m+1} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-m+1} \cdots p_{n+1}}{p_{n-m-1}}\right) \\ &\quad + (p_{n-m} - 1) \frac{p_{n-m} \cdots p_{n+1}}{p_{n-m}} \end{aligned}$$

and

$$\begin{aligned} g\left(\frac{p_{i_1} \cdots p_{i_{m+2}}}{p_{i_{m+2}}}, \dots, \frac{p_{i_1} \cdots p_{i_{m+2}}}{p_{i_1}}\right) &= p_{i_1} g\left(\frac{p_{i_2} \cdots p_{i_{m+2}}}{p_{i_{m+2}}}, \dots, \frac{p_{i_2} \cdots p_{i_{m+2}}}{p_{i_2}}\right) \\ &\quad + (p_{i_1} - 1) \frac{p_{i_1} \cdots p_{i_{m+2}}}{p_{i_1}} \end{aligned}$$

By induction, we obtain

$$g\left(\frac{p_{n-m+1} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-m+1} \cdots p_{n+1}}{p_{n-m-1}}\right) \geq g\left(\frac{p_{i_2} \cdots p_{i_{m+2}}}{p_{i_{m+2}}}, \dots, \frac{p_{i_2} \cdots p_{i_{m+2}}}{p_{i_2}}\right)$$

Moreover, since $i_1 < \dots < i_{m+2} \leq n+1$ then $i_{m+2-j} \leq n+1-j$ for each $j = 0, \dots, m+1$ and thus $p_{i_{m+1-j}} \leq p_{n-j}$. Therefore, $p_{i_1} = p_{i_{m+2}-(m+1)} \leq p_{n+1-(m+1)} = p_{n-m}$ and

$$\begin{aligned}
\frac{p_{n-m} \cdots p_{n+1}}{p_{n-m}} &= p_{n-m+1} p_{n-m+2} \cdots p_{n+1} \\
&= p_{n+1-m} p_{n+1-(m-1)} \cdots p_{n+1} \\
&\geq p_{i_{m+2}-m} p_{i_{m+2}-(m-1)} \cdots p_{i_{m+2}} \\
&= p_{i_2} p_{i_3} \cdots p_{i_{m+2}} \\
&= \frac{p_{i_1} \cdots p_{i_{m+2}}}{p_{i_1}}
\end{aligned}$$

Obtaining that

$$g\left(\frac{p_{n-m} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-m} \cdots p_{n+1}}{p_{n-m}}\right) \geq g\left(\frac{p_{i_1} \cdots p_{i_{m+2}}}{p_{i_{m+2}}}, \dots, \frac{p_{i_1} \cdots p_{i_{m+2}}}{p_{i_1}}\right).$$

□

We may now prove Theorem 1.4.

Proof of Theorem 1.4. Let $1 < p_1 < \cdots < p_{n+1}$ be prime integers, $n \geq 1$. We consider Theorem 2.1, where, for each $i = 1, \dots, n+1$, we let $x_i^j = p_i$ for all $1 \leq j \leq n$. Then, $(\underbrace{a \times \cdots \times a}_n)$ can be tiled with $(\underbrace{p_1 \times \cdots \times p_1}_n), \dots, (\underbrace{p_{n+1} \times \cdots \times p_{n+1}}_n)$ if

$$a > g_n = \max_{1 \leq k \leq n} \{g(X_k(i_1, \dots, i_{k+1})) | \{i_1, \dots, i_{k+1}\} \subseteq \{1, \dots, n+1\}\}.$$

By Lemma 2.3 (b), for each fixed $k = 1, \dots, n$

$$\begin{aligned}
\bar{g}_k &= \max\{g(X_k(i_1, \dots, i_{k+1})) | \{i_1, \dots, i_{k+1}\} \subseteq \{1, \dots, n+1\}\} \\
&= g\left(\frac{p_{n-k+1} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-k+1} \cdots p_{n+1}}{p_{n-k+1}}\right)
\end{aligned}$$

Moreover, for each $k = 1, \dots, n$, $\bar{g}_{k+1} > \bar{g}_k$ since

$$\begin{aligned}
\bar{g}_{k+1} &= g\left(\frac{p_{n-k} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-k} \cdots p_{n+1}}{p_{n-k}}\right) = p_{n-k} g\left(\frac{p_{n-k+1} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-k+1} \cdots p_{n+1}}{p_{n-k-1}}\right) \\
&\quad + (p_{n-k} - 1) \frac{p_{n-k} \cdots p_{n+1}}{p_{n-k}} \\
&> g\left(\frac{p_{n-k+1} \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_{n-k+1} \cdots p_{n+1}}{p_{n-k-1}}\right) = \bar{g}_k
\end{aligned}$$

So, $g_n = \max_{k=1, \dots, n} \{\bar{g}_k\}$ and, by Lemma 2.3 (a),

$$g_n = g\left(\frac{p_1 \cdots p_{n+1}}{p_{n+1}}, \dots, \frac{p_1 \cdots p_{n+1}}{p_1}\right) = n \prod_{i=1}^{n+1} p_i - \sum_{i=1}^{n+1} \frac{p_1 \cdots p_{n+1}}{p_i}.$$

□

Proposition 2.4 Let L, a, b, c and r be positive integers with $b|r$ and such that $r = x_1 a + x_2 c$ for some integers $x_1, x_2 \geq 0$ and $Lc = y_1 a + y_2 b$ for some integers $y_1, y_2 \geq 0$. Then, $(r + ac \times r + ac)$ and $(Lc + kab \times Lc + kab)$ can be tiled with $(a \times a)$, $(b \times b)$ and $(c \times c)$ for any integer $k \geq 1$.

Proof. Suppose that $b|r$. By Theorem 1.1, we have that

- $(r \times r)$ can be tiled with $(b \times b)$
- $(ac \times ac)$ can be tiled with $(a \times a)$,
- $(ac \times r)$ can be tiled with $(a \times a)$ and $(c \times c)$,
- $(Lc \times Lc)$ can be tiled with $(c \times c)$,
- $(Lc \times kab)$ can be tiled with $(a \times a)$ and $(b \times b)$ and
- $(kab \times kab)$ can be tiled with $(a \times a)$ (or $(b \times b)$).

The results follow by sticking together copies of the above rectangles as shown in Figure 1. □




Figure 1: Compositions of tilings

Proof of Theorem 1.5. By Theorem 1.1, we have that $(a \times a)$ can be tiled with (2×2) and (3×3) if $a \equiv 0 \pmod{2}$ or $a \equiv 0 \pmod{3}$. So, we only need to show that $(a \times a)$ can be tiled with (2×2) , (3×3) and $(p \times p)$ for any odd integer $a \geq 3p + 2$ with $a \equiv 1$ or $2 \pmod{3}$. We have two cases.

Case A) $p \equiv 1 \pmod{3}$. Let $s = p - 1 + k3 \geq p + 2$ for any integer $k \geq 1$. Since $s > g(2, p) = p - 1$ then there exist nonnegative integers u and v such that $s = 2u + pv$. Since $3|s$ then, by Proposition 2.4, $(a \times a) = (s + 2p \times s + 2p)$ can be tiled with (2×2) , (3×3) and $(p \times p)$ for any $a = s + 2p \geq p + 2 + 2p = 3p + 2$ and $a \equiv 2 \pmod{3}$.

Now, since $p = 3k + 1$ for some integer $k \geq 1$ then for $p > 3$ we have that $p = (k - 1)3 + 2(2)$ and by Proposition 2.4 (with $L = 1$), we have that $(a \times a) = (p + k6 \times p + k6)$ can be tiled with (2×2) , (3×3) and $(p \times p)$ for any odd integer $a = p + k6 \geq p + 6$ with $a \equiv 1 \pmod{3}$.

Case B) $p \equiv 2 \pmod{3}$. Let $s = p - 2 + k3 \geq p + 1$ for any integer $k \geq 1$. Since $s > g(2, p) = p - 1$ then there exist nonnegative integers u and v such that $s = 2u + pv$. Since $3|s$ then, by Proposition 2.4, $(a \times a) = (s + 2p \times s + 2p)$ can be tiled with (2×2) , (3×3) and $(p \times p)$ for any $a = s + 2p \geq 2p + 1 + p = 3p + 1$ and $a \equiv 1 \pmod{3}$.

Now, since $p = 3k + 2$ for some integer $k \geq 1$ then for $p > 3$ we have that $p = (k - 1)3 + 2(2)$ and by Proposition 2.4 (with $L = 1$), we have that $(a \times a) = (p + k6 \times p + k6)$ can be tiled with (2×2) , (3×3) and $(p \times p)$ for any odd integer $a = p + k6 \geq p + 6$ with $a \equiv 2 \pmod{3}$.

Let us set $p = 5$. It is clear that (1×1) and (7×7) cannot be tiled with (2×2) , (3×3) and (5×5) . By the above cases, we have that $(a \times a)$ can be tiled with (2×2) , (3×3) and (5×5) if $a \geq 3p + 2 = 17$ and, by Theorem 1.1, $(a \times a)$ can be tiled with (2×2) and (3×3) if $a \equiv 0 \pmod{2}$ or $a \equiv 0 \pmod{3}$. This leave us the cases when $a = 5, 11$ and 13 . If $a = 5$ is trivial. (11×11) can be tiled with (2×2) , (3×3) and (5×5) since, by tha above case B, the result is true for any odd integer $a \geq p + 6 = 11$ and $a \equiv 2 \pmod{3}$. Finally, (13×13) can be tiled as it is illustrated in Figure 2.

5		2	2	2	2
		2	2	2	2
		2	2	2	2
2	3	2	2	2	2
2		2	2	2	2
2	5		2	2	2
2			3	3	

Figure 2: Tiling of (13×13) with bricks (2×2) , (3×3) and (5×5)

Let us set $p = 7$. It is clear that (1×1) , (5×5) and (11×11) cannot be tiled with (2×2) , (3×3) and (7×7) . By the above cases, we have that $(a \times a)$ can be tiled with (2×2) , (3×3) and (7×7) if $a \geq 3p + 2 = 23$ and, by Theorem 1.1, $(a \times a)$ can be tiled with (2×2) and (3×3) if $a \equiv 0 \pmod{2}$ or $a \equiv 0 \pmod{3}$. This leave us the cases when $a = 7, 13, 17$ and 19 . If $a = 7$ is trivial. (13×13) and (19×19) both can be tiled since, by the above case A, the result is true for any odd integer $a \geq p + 6 = 13$ with $a \equiv 1 \pmod{3}$. Finally, (17×17) can be tiled as it is illustrated in Figure 3.

7			2	2	2	2	2
			2	2	2	2	2
			2	2	2	2	2
			2	2	2	2	2
2	2	3	2	2	2	2	2
2	2		2	2	2	2	2
2	2	7			2	2	2
2	2				2	2	2
2	2				3	3	

Figure 3: Tiling of (17×17) with bricks (2×2) , (3×3) and (7×7)

□

Acknowledgement

I would like to thank Y. O. Hamidoune for several helpful conversations.

References

- [1] F.W. Barnes, Algebraic theory of bricks packing II, *Discrete Mathematics* **42** (1982), 129-144.
- [2] A. Brauer and J.E. Shockley, On a problem of Frobenius, *J. Reine Angewandte Math.* **211** (1962), 215-220.
- [3] R.J. Brower and T.S. Michael, When can you tile a box with translates of two given rectangular bricks?, *Elec. J. of Comb.* **11** (2004), # N7.
- [4] N.G. de Bruijn, Filling boxes with bricks, *Amer. Math. Monthly* **76** (1969), 37-40.
- [5] J. Fricke, Quadratzerlegung eines Rechtecks, *Math. Semesterber.* **42** (1995), 53-62.
- [6] S.M. Johnson, A linear diophantine problem, *Can. J. Math.* **12** (1960), 390-398.
- [7] D.A. Klarner, Packing a rectangle with congruent n -ominoes, *J. Comb. Theory* **7** (1969), 107-115.
- [8] L.F. Klosinski, G.L. Alexanderson and L.C. Larson, The Fifty-Second William Lowell Putman Mathematical Competition, *Amer. Math. Month.* **9** (1992), 715-724.
- [9] D. A. Narayan and A.J. Schwenk, Tiling large rectangles *Mathematics Magazine* **75**(5) (2002), 372-380.
- [10] T. Plambeck, Web site <http://www.plambeck.org/oldhtml/mathematics/klarner/>
- [11] J.L. Ramírez Alfonsín, The Diophantine Frobenius Problem, *Oxford Lectures Series in Mathematics and its Applications* **30**, Oxford University Press, (2005).